

T
U
C
K

T
O
T
A
L

P
A
U

20

J
U
N
E
2
0
0
8

Valorisation du CO₂ par l'industrie chimique


CO₂ as building block in the chemical industry

Michele Aresta

CIRCC and Department of Chemistry
University of Bari, Campus Universitario
70126 Bari, Italy

m.aresta@chimica.uniba.it


Where we are....



Outline

- The Carbon Cycle
 - The anthropogenic carbon dioxide
- The Origin of Anthropogenic CO₂
 - How we can use CO₂
- Industrial Utilisation of CO₂
 - **Chemical**
 - **Technological**
- The Must in the Utilisation of CO₂
 - Used vs avoided CO₂
- Sustainability and CO₂ Utilisation
 - Process innovation
 - Tandem chemical-biological
 - Conversion of biomass
- **Classes of Compounds of Industrial Interest**
- LCA as a Tool for Quantifying the Avoided CO₂
- Perspective in CO₂ Utilisation
 - Opportunities and Challenges

The Carbon Cycle and the Excess Anthropogenic CO₂



The origin of anthropogenic CO₂

**90 % Use of fossil carbon
and biomass for energy
production**

10 % from chemicals

How can we use CO₂ ?

Fuels:

HCs require hydrogen
De-oxygenation to C
requires energy

Chemicals:

Bulk chemicals
Specialties

The Utilization of CO₂ in the Chemical Industry

Compound	Date	Amount Mt/y	Catalyst
Urea	1870	70	No cat
Salycilic acid	1869	20 E-3	Group 1 metals
Methanol	1970	6	TM oxides
Carbonates			
Organic	1972	Few kt	Various systems
Inorganic	1860	30	No cat

The Technological Utilization

18 Mt/y

- Additive to beverages
- Food packing/conservation
- Cereals preservation (bactericide)
- Fire extinguisher
- Water treatment
- Mechanical industry (moulding,...)
- Dry-washing
- Air-conditioning
- Extraction (scf and EOR)

The MUST in the utilisation of
 CO_2

The new process (conversion
or technological use) or
product (substitute of existing
ones) must minimize
the energy consumption
and the emission of
 CO_2

Used CO₂ *versus* Avoided CO₂





- **The lifetime of chemicals**
 - Storage of carbon: limited potential**
- **Recycling of carbon**
 - Innovative processes**
- **Rate of formation and fixation**
 - Combustion processes vs conversion**
- **Reducing fossil-C extraction**
 - Efficient conversion processes**
 - Optimized C-utilization efficiency**

Avoid the extraction of CO₂

- Use of recovered CO₂
 - Purity (food industry)
 - Better exploitation of sources of pure CO₂
 - Fermentation industry
 - » Seasonality
- Replace extracted CO₂
 - EOR

Sustainability and CO₂ Utilisation

- **Process Innovation**
(use of concentrated CO₂-flows):
 - Reduce the Carbon and Energy dependence
 - Recycling Carbon
 - Raw material diversification
- **Tandem Chemical-Biological**
 - Coupling Chemistry and Biotechnology
- **Intelligent use of Biomass**
(exploitation of atmospheric CO₂)
 - Nature makes, Man re-shapes



A

B

D

C


H₂

e⁻, H⁺

Classes of compounds of industrial interest: opportunities and challenges

- **Carboxylates:**
 - Formic acid
 - Long chain acids, lactones, pyrones
- **Carbonates**
 - Molecular (linear and cyclic) and polymeric
- **Carbamates**
 - Molecular and polymeric
- **Acrylic acids and their esters**
- **Energy products**


Synthesis of Formic Acid



Ru

Rh

Leitner et al., *J. Chem. Soc. Chem. Commun.*, 1465-1466, 1993



Amines




Jessop et al., *Nature*, 368, 231-233, 1994

TOFh ~ 10000 at 80 °C and 1364 at room temperature

High pressure: Laurency et al. *Appl. Catal. A: Gen.*, 2003; *Chem. Eur. J.* 2007

Lactones, Pyrones, Long Chain Carboxylic Acids or Esters



Musco, 1978; Behr, 1984; Braunstein, 1998; Dinjus, 2002


M.Aresta, New J. Chem, 1994, 18, 133

Carboxylation of Alcohols



- **Thermodynamics** says that the reaction is “quasi-neutral”: the equilibrium concentration of the carbonate is reported to be close to 1-2%
- **Water** is formed that should be eliminated for displacing the equilibrium to right and avoid that catalysts are modified
- The **reaction mechanism**: in principle an alcohol should undergo an **acid-activation** (to generate **-R**) and a **base-activation** (to generate **-OR**): bi-functional catalysis


Double base-activation of methanol



Transition metal system



T. Sakakura, 1998-2000; D. Ballivet, 2002





Aresta et al., 2003, 2005, 2006, 2007



Reaction mechanism of formation of DMC from methanol and CO₂ under [Nb(OCH₃)₅]₂ catalysis: the case of the base-plus-acid activation of methanol

Use of heterogenized catalysts

Grafted Sn-complexes



Catalyst recycled ten times with a DMC/Sn ratio equal to 1.2-1.6 per cycle




M. Aresta et al, ICA, 2007

Use of heterogeneous catalysts

CeO_2 (Tomishige 2006)	Short-living
ZrO_2 (Tomishige 1999)	Short-living
$\text{CeO}_2/\text{ZrO}_2$	(Tomishige 2000)
Tl_2CO_3 (Aresta, 2006)	Long-lasting, slow
$\text{CeO}_2\text{-Al}_2\text{O}_3$ (Aresta, 2008)	Long-lasting, fast
$\text{CeO}_2 - \text{M}_x\text{O}_y$ (Aresta, 2008)	Long-lasting, fast

Effect of catalyst modification on its performance




Commercial CeO_2
Size of particles: μm

Each cycle 5 h

CeO_2 nano-particles
synthesised and
loaded with Al and Fe
Each cycle 3 h


Aresta et al, CatTod 2008



Cyclic carbonates

- Epoxides plus CO_2
 - Several catalysts are used
 - IL are optimal reaction media as they fasten the reaction
- Olefins plus CO_2 and O_2
(oxidative carboxylation)
 - One-pot synthesis avoiding the isolation of epoxides

Direct carbonation of olefins. Several issues.....



1. The oxidation products distribution is mediated by CO_2

2. An aldehyde is formed that promotes the formation of the epoxide

3. The latter is converted into the carbonate


4. Is it possible to avoid the double bond cleavage?

Detailed study on solvent, pressure of O_2 and CO_2 , co-catalysts

Carbonate formation without the intermediacy of the radical path. Surface chemistry on metal-oxides!


A two-step process based on molecular oxygen

- $\text{Nb}_v=\text{O} + \text{CO}_2 \rightarrow \text{NbCO}_3$ (*FTIR, literature*)
- Lattice-oxygen (M_yO_x) can be transferred to the olefin \rightarrow “ $\text{O}_{x-1}\text{M}_y\text{OCH}_2\text{CHPh}$ ”. (*CatTod, 2006; CatTod, 2008*)
- NbCO_3 can react with oxidized styrene \rightarrow Nb(III) + carbonate




Asymmetric syntheses

Enantiomer R



Enantiomer S



R	Catalyst	Yield*	TON	e.e.
CH ₃	Nb ₂ O ₅	77 %	17	> 98% (R)
Ph	Nb ₂ O ₅	80 %	29	> 98% (R)





R	Catalyst	Yield*	TON	e.e.
CH ₃	Nb ₂ O ₅	65 %	15	> 98% (S)
Ph	Nb ₂ O ₅	65 %	25	> 98% (S)

* Isolated yield

CO₂ as co-monomer

- Co-polymerisation with propene-epoxide
 - Al-porphyrins (on stream in CH)
- Co-polymerisation with cyclohexene-oxide
 - Cr-catalysts
 - Zn-catalysts


Coupling of ethene with CO₂




The elimination of acrylic acid is not observed although the thermodynamics is favourable: The kinetic barrier is too high. Energy of the M-O and M-H bonds

Avoiding the formation of the M-H bond favours the elimination of the acrylate: the mechanism of formation of the ester from CO₂ and ethene is still under investigation.


Aresta et al, Chem Eu J, 2007



The carboxylation of glycerol: a valuable product from two wastes





M. Aresta, A. Dibenedetto, F. Nocito, C. Pastore, *J. Mol. Catal.: A*, 2006




Catalyst fully recovered and reused; 90 % isolated in 2 h at 140 °C
Molar ratio Urea : Glycerol = 1:1 weight ratio Catalyst : Glycerol 0.03

M. Aresta, A. Dibenedetto, F. Nocito, Patent Application

CO₂ as mild oxidant



Methane Tri-Reforming Plasma technology




The methane dry-reforming may contribute to the implementation of the GTL technology with a direct conversion at the extraction well of LNG into liquid fuels

Coupling Chemistry and Biotechnology: From CO₂ and H₂O to CH₃OH at rt

- $\text{CO}_2_{\text{aq}} \rightarrow \text{HCOO}^-$ Formate dehydrogenase
- $\text{HCOO}^- \rightarrow \text{H}_2\text{CO}$ Formaldehydedehydrogenase
- $\text{H}_2\text{CO} \rightarrow \text{CH}_3\text{OH}$ Methanoldehydrogenase
- **NAD⁺/NADH is the source of energy.**
- **Key issue: how to reverse the NAD⁺/NADH couple after oxidation?**
- **Use of TM systems for solar light harvesting-utilisation**

Enzymatic Phenol Carboxylation

Biotechnological synthesis of 4-OH benzoic acid
also in *sc*-CO₂



REAGENT
+
COSUBSTRATES
+
ENZYME

*Inglobated into
Low-melting
agar*




PRODUCT

Aresta, M., Tommasi, I., Dileo, C.
Tetrahedron, 54(1998), 8841 - 8846

Reagent
+
Cosubstrates
+
Enzymes

*Membrane
reactor*



Product

Aresta M., Dibenedetto A. *Rev.
Mol. Biotech.*, 90, 113-128, 2002

Biomass exploitation:

Nature makes, Man re-shapes

Non-destructive

- Extraction of chemicals with complex molecular structure, including triglycerids for biodiesel (Product Entropy)



Partially destructive

- Exploitation of the lignine structural features
- Enzymatic conversion of lignine and cellulosic materials

Destructive

- Biogas production
- Conversion of biomass into Syngas
- Use of Syngas for the synthesis of methanol
 - Bio-methanol for FAME
 - Bio-hydrogen (bacterial or chemical production)
- Use of Syngas for the synthesis of gasoline and diesel

Quantifying the avoided CO₂



Opportunities and challenges over 1 Mt/y

- Carbonates
- Acrylic acids and their esters
- Fuels
- Polymers
- Olefins oxidative carboxylation
- Water elimination
- Avoid the formation of hydrido-acrylates
- Source of hydrogen
 - Water splitting
 - Reduction of CO₂ in water
- Production of epoxides and azetidines/aziridines
- Suppression of the radical path



Catalyst life

